

COCHRANE UPDATED REVIEW

Endovascular Ruptured Abdominal Aortic Aneurysm Repair (EVRAR): A Systematic Review[☆]

D.W. Harkin,^{1*} M. Dillon,¹ P.H. Blair,¹ P.K. Ellis¹ and F. Kee²

¹Regional Vascular Surgery Unit, Royal Victoria Hospital, Belfast BT12 6BA, Northern Ireland, UK, and

²Department of Epidemiology and Public Health, Queen's University Belfast, University Road, Belfast BT7 1NN, Northern Ireland, UK

Background. To review evidence supporting the use of endovascular ruptured aneurysm repair (EVRAR) for treatment of ruptured abdominal aortic aneurysm (RAAA).

Methods. A systematic review of the medical literature was performed for relevant studies. We searched a number of electronic databases and hand-searched relevant journals until November 2006 to identify studies for inclusion. We considered studies in which patients with a confirmed ruptured abdominal aortic aneurysm were treated with EVRAR, which reported endpoints of mortality and major complications.

Results. There was 1 randomised controlled trial (RCT), 33 non-randomised case series (24 retrospective and 9 prospective) reports were identified comparing EVRAR (n = 891) with conventional open surgical repair for the treatment of RAAA. Whilst no benefit in the primary outcome of mortality was noted in the only RCT, evidence from non-randomised studies suggest that EVRAR is feasible in selected patients, where it may be associated with a trend towards reductions in blood loss, duration of intensive care treatment, early complications, and mortality.

Conclusions. For the treatment of symptomatic or ruptured abdominal aortic aneurysm, emergency endovascular repair (EVRAR) is feasible in selected patients, with early outcomes comparable to best conventional open surgical repair for the treatment of RAAA.

© 2007 European Society for Vascular Surgery. Published by Elsevier Ltd. All rights reserved.

Keywords: Abdominal aortic aneurysm; Rupture; Surgical treatment; Endovascular aneurysm repair (EVAR); Endovascular ruptured aneurysm repair (EVRAR); Systematic review.

Introduction

Abdominal aortic aneurysm (AAA), affects between 1.2% and 7.6% of the population over 50 years of age in the United Kingdom.^{1,2} Untreated, progressive aneurysm enlargement can lead to rupture, which is lethal in 80% of those affected and responsible for over 6800 deaths per annum in the United Kingdom and 2.1% of all deaths in men over 65 years.^{2,3} Excellent results can now be achieved with elective open repair, with some specialist centres reporting mortality rates of less than 2% and surgeons in non-specialist units achieving mortality rates of 5% to 8%.^{4–7} This contrasts with

emergency open repair for a ruptured AAA (RAAA), in these critically ill patients, which carries a mortality rate of 30% to 65%.^{4,8,9} In the last two decades a new minimally-invasive technique, endovascular aneurysm repair (EVAR), has offered an alternative therapy to conventional open repair for selected patients with AAA, and has shown significant reductions in early complications and mortality.^{10,11} Furthermore, several studies have shown that EVAR, especially under local anaesthesia, reduces the physiological insult to the body as compared to conventional open surgical repair.^{12,13} This has led some to consider emergency endovascular aneurysm repair (EVRAR) as an alternative to conventional open repair in selected patients with RAAA.^{13–15}

This article reviews the available published evidence to support the use of emergency endovascular aneurysm repair (EVRAR) as compared to conventional open repair (OR) of RAAA, for patients with RAAA.

[☆] Update of Cochrane Review: Cochrane Database Syst. Rev 2006: (1);CD005261

*Corresponding author. D. W. Harkin, MD FRCS EBSQ-VASC, Regional Vascular Surgery Unit, Royal Victoria Hospital, Grosvenor Road, Belfast BT12 6BA, Northern Ireland, UK.

E-mail address: denis.harkin@royalhospitals.n-i.nhs.uk

Methods

Criteria for considering studies

Our objective was to review the published evidence to allow assessment of the advantages and disadvantages of EVAR for patients with RAAA (EVRAR). This was determined by the effect on short-term mortality, major complication rates, aneurysm exclusion, and late complications when compared with patients who have had conventional open repair of RAAA. To ensure comparability of participants were considered for inclusion. There must have been evidence of rupture on imaging, computerised tomography angiography (CTA) or magnetic resonance angiography (MRA), or objective acute symptoms suggestive of impending rupture of a known aneurysm (abdominal or back pain in a patient with an aneurysm) to warrant inclusion. Studies where objective evidence of RAAA is not clear were excluded.

We did not limit our search in respect to the type of endovascular intervention or device type.

The following outcome measures were considered: (1) mortality (30 day, or in-hospital mortality, i.e. procedure related); (2) aneurysm exclusion (no flow in the AAA sac, or further extravasation (escape of blood from the vessel into the tissues) beyond the sac on follow-up imaging 30 days after the procedure); (3) comparative surrogate measures of procedural success (blood loss, procedural time, ICU stay). Data was sought for other important outcome measures, such as: major complications (i.e. open conversion, haemorrhage, myocardial infarction, stroke, renal failure, respiratory failure, pneumonia, bowel ischaemia, lower limb ischaemia); minor complications (i.e. haematoma, wound infection); long term complications, re-intervention rates and mortality. However, these data were inconsistently reported and as such comments were possible pertain to individual studies.

Search strategy

We searched for articles using the broad terms "endovascular" AND "abdominal aortic aneurysm" AND "ruptured" in all databases. The last search prior to preparation of this review was performed in November 2006. We searched the Cochrane Peripheral Vascular Diseases Group trials register (last searched October 2006) and the Cochrane Central Register of Controlled Trials (CENTRAL) database (last searched Issue 4, 2006) for trials describing endovascular repair of ruptured or symptomatic abdominal aortic aneurysm.

We performed electronic searches of the following bibliographic databases: (1) AMED (Allied and Complementary Medicine Database); (2) Best Evidence; (3) Biological Abstracts (4) HMIC (Health Management of Information Consortium – comprising DH-data, the King's Fund Database and Helmis); (5) NHS DARE (Database of Assessments of Reviews of Effects); (6) NHS EED (Economics Evaluations Database); (7) NHS HTA (Health Technology Assessment); (8) PubMed; (9) Science Citation Index; (10) MEDLINE. In order to collect any articles missed by searches we also hand-searched relevant journals, retrieved articles, and relevant. Searches were not restricted by publication type, study design or language of publication.

Selection

Authors, M.D. and D.W.H., evaluated the trials under consideration independently for appropriateness for inclusion and for methodological quality. Disagreements were resolved on discussion with the review team and agreed arbitrators. The searches identified 41 potentially relevant articles comparing EVRAR with conventional open surgical repair for the treatment of RAAA to which exclusion criteria were applied: 1 randomised controlled trial; 33 case series (24 retrospective and 9 prospective); 7 case reports.

Only 1 RCT was identified, and as such no tests of heterogeneity or sensitivity analysis were performed, and meta-analysis was not indicated.

Results

There was only 1 completed randomised controlled trial (RCT)¹⁶ comparing EVRAR with conventional open surgical repair for the treatment of RAAA. Thirty-three case series (24 retrospective and 9 prospective)^{13–15,17–46} describing EVRAR (in 876 patients) were identified, [Table 1](#). A number of case-control studies and cohort studies comparing outcomes of EVRAR with open surgical repair (OR) were found,^{13,15,17–19,22,24,27,30,31,35,38,41,44–46} these were invariably non-randomised and in some cases the EVRAR had been compared to historical controls. Therefore, the following review has limited high-quality evidence on which to base strong recommendations, but endeavours to present the available evidence as succinctly as possible.

Hinchcliffe *et al.*,¹⁶ in late 2006 reported the results of a pilot study as part of an ongoing randomised controlled trial comparing endovascular (EVRAR) and open surgery (OR) for the treatment of ruptured abdominal aortic aneurysm (RAAA). The trialists

Table 1. Mortality after endovascular ruptured aneurysm repair (EVRAR)

Study	Design	%EVAR Suitable*	EVRAR	OPEN group (if quoted)	<i>p</i> value (if quoted)
Hinchcliffe 2006	PRCT	93%	8/15 (53)	9/17 (53%)	NS
Franks 2006	PCS†	100%	1/10 (10)	7/10 (70)	—
Arya 2006	PCCS	42%	4/17 (24)	11/23 (47)	NS
Lagana 2006	CS		4/38 (10.5)		
Visser 2006	RCS	47%	8/26 (31)	9/29 (31)	NS
Alsac 2005	PCS	73%	4/17 (23.5)	10/20 (50)	0.09
Brandt 2005	RCS	72%	0/11 (0)	2/13 (15)	NS
Greco 2005	RCS	—	114/290 (39)	2627/5508 (47)	0.05
Kapma 2005	RCS	36%	5/40 (13)	64/213 (30)	0.02
Larzon 2005	RCS	—	2/15 (13)	12/26 (46)	0.05
Peppelenbosch 2005	RCS		8/35 (23)		—
Vaddineni 2005	RCS	60%	2/9 (22)	4/15 (26)	NS
Castelli 2005	CS		5/25 (20)		
Gerassimidis 2005	CS		9/23 (39)		
Hechelhammer 2005	CS		4/37 (10.4)		
Mehta 2005	CS		7/30 (23)		
Lombardi 2004	CS		0/5 (0)		
Lee 2004	RCS	88%	1/13 (7.7)	1/4 (25)	—
Rubin 2004	CS		1/5 (20)		
Peppelenbosch 2003	PCS†	80%	4/26 (15)	4/14 (28)	—
Reichart 2003	PCS	42%	1/6 (16.6)	4/13 (30)	—
Resch 2003	PCS	79%	4/14 (29)	8/23 (35)	0.05
Scharrer-Pamler 2003	CS		3/24 (12.5)		
Van Herzeele 2003	CS		1/9 (11)		
Verhoeven 2002	PCS	34%	1/9 (11)	7/31 (23)	—
Yilmaz 2002	PCS	81%	4/17 (24)	12/29 (41)	NS
Doss 2002	CS		0/6 (0)		
Lachat 2002	CS		2/21 (9.5)		
Orend 2002	CS		4/21 (19)		
Van Sambeek 2002	CS		6/22 (19)		
Hinchcliffe 2001	CS		9/20 (45)		
Ohki 2000	PCS	80%	2/20 (10)	0/5 (0)	—
Greenberg 2000	CS		0/3 (0)		
Ohki 1999	CS		2/12 (17)		
Overall ^{13,15,16,18–47}		67% (34–100)	18% (0–53)	34% (0–70)	

Data in parenthesis represents (%) or (range).^{13,15,16,18–47} Prospective Randomised Controlled Trial (PRCT); Prospective case-control study (PCCS); Prospective Cohort Study (PCS); Retrospective Cohort Study (RCS); Case series (CS). *Percentage of cohort considered for endovascular repair deemed anatomically suitable after contrast-enhanced CT scan. †Control group historical.

considering 103 patients with suspected ruptured AAA, and randomised 32 patients (15 to EVRAR and 17 to OR). This low rate of patient randomisation (31%) may be in part explained by the broad exclusion criteria applied, including: patient unfit for open repair; death before randomisation; inability to give consent; age; refusal of operation; no team available; surgeon preference. This trial highlights the significant logistical problems related in having a fully trained, equipped, and committed endovascular team available at all times. On an intention to treat basis there was no significant difference in mortality for the EVRAR group, 8 of 15 (53%) compared to OR group, 9 of 17 (53%). If we exclude those patients who died after randomisation but before intervention, mortality for the EVRAR group, 7 of 13 (54%) including 2 EVRAR failures requiring open conversions (1 survivor; 1 non-survivor), compared to OR group, 6 of 14 (43%), non-significant. Of patients who survived intervention, there was no significant difference in postoperative

complications (moderate and severe) for the EVRAR group, 77%, compared to OR group, 80%. Severe renal complications were significantly more common in the EVRAR group, 6 of 11 (55%), compared to OR group, 1 of 14 (7%), $p < 0.02$. A significant difference in recorded blood loss and blood transfusion requirements is recorded but difficult to interpret when comparing minimally invasive, in which intra-abdominal blood loss cannot be counted, and open surgical procedures. Whilst the authors should be complemented on their enthusiasm to answer this question, their results must be considered with caution due to logistical deficiencies in the availability of a 24-hour endovascular service at their unit, subjective entry criteria, and failure to recruit sufficient numbers.

The results of non-randomised case series comparing EVRAR, to contemporary or historical series of open repair (OR) for the treatment of RAAA,^{13,15,17–19,22,24,27,30,31,35,38,41,44–46} are compared to the 1 available RCT¹⁶ and presented in Tables 1–4.

Table 2. Length of ICU stay

Study	eEVAR group	OPEN group	<i>p</i> value (if quoted)
Franks 2006	1.3	6.1	0.01
Alsac 2005	3	13	0.01
Brandt 2005	4.8	8.5	NS
Kapma 2005	0*†	48*†	0.001
Vaddenini 2005	5*	20*	—
Peppelenbosch 2003	46*†	154*†	—
Reichart 2003	2.25	13 days	—
Resch 2003	1*	3*	0.02
van Sambeek 2002	8*†	62*†	0.004
Yilmaz 2002	2.2	5.2	0.05

Data represents mean unless indicated *median. Values in days except were indicated †hours.

Discussion

Abdominal aortic aneurysm

Abdominal aortic aneurysm (AAA), affects between 1.2% and 7.6% of the population over 50 years of age in the United Kingdom. The prevalence of AAA in men is approximately three times greater than in women, and the incidence increases with advancing age.^{1,2} The risk of aneurysm rupture has been shown to be proportional to aneurysm size, with aneurysms measuring less than 5.0 cm having an annual rupture rate of approximately 1% whereas those greater than 7.0 cm in diameter have an annual rupture rate of over 30%.^{5,47} The UK Small Aneurysm Trial has shown that in general, patients benefit from aneurysm repair when maximum aneurysm diameter exceeds 5.5 cm, at which stage the risk of spontaneous rupture exceeds the risks of conventional open surgical repair.⁵ Excellent results can now be achieved with conventional open repair, with some specialist centres reporting mortality rates of less than 2% and surgeons in non-specialist units achieving mortality rates of 5% to 8%.^{5,7,48} Furthermore, a newer minimally-invasive technique called endovascular aneurysm repair

Table 3. Length of procedure

Study	EVRAR group	OPEN group	<i>p</i> value (if quoted)
Hinchcliffe 2006	160	150	0.34
Franks 2006	156	186	0.04
Alsac 2005	156	222	0.1
Brandt 2005	178	207	NS
Kapma 2005	110*	180*	0.001
Vaddenini 2005	143	181	—
Peppelenbosch 2003	154	155	—
Verhoeven 2002	110	122.5	—
van Sambeek 2002	193	203	NS
Yilmaz 2002	173	273	0.05
Ohki 2000	336*	492*	—

Data represents mean unless indicated *median. Values in minutes.

Table 4. Blood loss and transfusion requirement

Study	EVRAR group	OPEN group	<i>p</i> value (if quoted)
A: Blood loss			
Hinchcliffe 2006	200	2100	0.004
Kapma 2005	200*	3500	0.001
Vaddenini 2005	475	2880	0.0001
Peppelenbosch 2003	1100	2600	—
Reichart 2003	300	4500	—
Resch 2003	800	4000	0.0001
van Sambeek 2002	125*	3400	0.01
Yilmaz 2002	660	3550	0.05
Ohki 2000	400*	2000	—
B: Transfusion requirements			
Hinchcliffe 2006	3	6	0.02
Franks 2006	0.86†	10.7†	0.01
Alsac 2005	1520	3075	0.1
Brandt 2005	964	1986	0.02
Kapma 2005	0*†	6†	0.001
Vaddenini 2005	3.78†	6.93†	0.014
Reichart 2003	0	1600	—
Resch 2003	2†	9†	0.02
Ohki 2000	3*†	6†	—

A: Data represents mean unless indicated *median. Values in millilitres. B: Data represents mean except were indicated *median. Values in millilitres except were indicated †units.

(EVAR), first reported by Parodi in 1991,⁴⁹ now offers an alternative treatment in certain patients. Two recent large prospective randomised controlled trials have compared elective EVAR with conventional open repair for the treatment of large AAA, and have shown significant reductions in early complications and mortality.^{11,50} However, these trials have also reinforced the knowledge that open repair is a successful technique and will remain a common form of treatment for over half of those patients presenting for whom EVAR is unsuitable on anatomical grounds or due to other factors.^{11,50} It is also clear from these studies that EVAR is associated with a higher re-intervention rate than open repair,^{11,50} and registry data would suggest that these re-intervention rates can remain constant and may even increase with time.⁵¹ As such long-term surveillance is essential after EVAR to monitor for endoleaks and stent integrity in order to reduce the small but significant incidence of late aneurysm rupture (Fig. 1).⁵²

Ruptured abdominal aortic aneurysm

Currently, rupture leads to death in over 80% of those affected, including 30% to 65% of those who receive conventional open surgical repair, and is responsible for over 6800 deaths per annum in the United Kingdom and 2.1% of all deaths in men over 65 years.^{2,4,9} The haemorrhagic shock and lower torso ischaemia-reperfusion injury which accompanies RAAA activate




Fig. 1. Algorithm for the treatment of patients presenting with ruptured abdominal aortic aneurysm. *EVAR suitable (see Table 5), †Hardman criteria.⁹

multiple inflammatory pathways in the body inducing a harmful proliferative systemic inflammatory response syndrome characterized by immune cells activation, pro-inflammatory mediator production and widespread vital organ injury (heart, lungs, liver, kidney, gut, etc.). The sequential failure of these organs despite intensive care support, once established leads to death in over 70% of cases.^{53–55} Detailed risk analysis and scoring systems have been shown to predict non-survivors in certain groups but individual patient outcomes cannot be accurately predicted.^{9,56} Clinicians have been reticent to apply these scoring systems rigidly as to do so would serve to preclude most patients with RAAA from surgical repair, condemning them to certain death.⁹ It is also now clear that those patients who undergo successful open repair of RAAA enjoy a post-operative quality of life similar to the “normal population”.⁵⁷

Endovascular ruptured aneurysm repair (EVRAR)

Since the first description of the EVAR technique,⁴⁹ many specialised vascular surgery centres have adopted its use in the elective treatment of abdominal aortic aneurysm, where its use has contributed to a reduction in early postoperative morbidity and mortality.¹¹ Furthermore, several studies have confirmed that the use of EVAR, especially under local anaesthesia, reduces the physiological insult to the body as compared to conventional open surgical repair.^{12,58} These potential benefits have led several experienced units to offer endovascular ruptured aneurysm repair (EVRAR), as an alternative to conventional open surgery, in selected patients.^{59,60} These early reports have suggested EVRAR is feasible and may be achieved with a morbidity and mortality in

selected patients which is at least equivalent to best conventional surgical practise.^{14,25,35}

Selection of cases on the basis of precise anatomical suitability for elective EVAR has been shown to be associated with a much lower rate of re-intervention, morbidity, and procedure-related cost.^{12,51,61} Evidence from several studies has shown that the aneurysm morphology is significantly more challenging for endovascular techniques in those assessed for RAAA compared with those undergoing elective EVAR.^{38,62,63} Most authors have used established criteria derived from elective EVAR to determine anatomical suitability for EVRAR. However, reported studies reveal substantial variation in the anatomical inclusion and exclusion criteria employed, with anatomical suitability rates ranging from 34% to 100%.^{15,16,22,31,38,44} In particular many groups accept inferior proximal neck anatomy which would preclude patients from elective EVAR, suggesting a trend to be more inclusive in these high risk patients.⁶⁴ Whilst there is no long term follow-up data available for patients undergoing EVRAR, evidence from elective EVAR would suggest that relaxation of the criteria for anatomical suitability may lead to future problems, such as increased rates of endoleak, graft displacement, complications, re-interventions or the need for open conversion.^{51,65}

The majority of centres use contrast-enhanced computed tomographic angiography (CTA) to assess the anatomical suitability of aneurysms for EVRAR.^{15,16,22,31,38,44,66–68} Some centres advocate intra-operative calibrated angiography as an effective alternative in order to reduce pre-operative delays.^{35,69} However, angiography does not clearly show luminal thrombus, which could adversely affect the ability to obtain a secure seal at the proximal fixation site risking endoleak or late device migration. Generous device oversizing is advocated and can improve the chances of a primary seal.^{64,65} However, concerns have been raised in relation to the effect of oversizing on aneurysm neck dilation and late failure.⁷⁰ Recent studies suggest CTA assessment can be safely carried out in rupture patients, suggesting any diagnostic delays may not be clinically important.^{66,71} Indeed the majority of patients with RAAA who are not operated upon survive for more than two hours after hospital admission and maintain a satisfactory systolic blood pressure greater than 80 mm Hg with minimal fluid resuscitation, which would allow sufficient time for radiological assessment in most specialist centres.^{66,71} In Table 5, we summarize current anatomical suitability criteria applied for emergency EVRAR in comparison to established criteria for elective EVAR.^{15,16,22,31,38,44,66–68}

Table 5. The Anatomical criteria used for consideration of EVRAR

Criteria	Elective EVAR	Emergency EVRAR	Comment
Neck diameter (max)	28 mm	32 mm	Allow for >10% oversize
Neck length (min)	15 mm	8 mm	Consider proximal fixation
Neck angulation (max)	60°	90°	Consider flexible device
Neck conicity (max)	4 mm/cm	8 mm/cm	Consider oversize
Neck quality	Moderate	Poor	Consider oversize
CIA/EIA diameter (min)	9 mm	6 mm	Consider uni-iliac device or conduit
CIA/EIA diameter (max)	16 mm	22 mm	Consider uni-iliac device
Iliac tortuosity	Moderate	Severe	Consider uni-iliac device or conduit

Data represents for elective EVAR average inclusion criteria for EVAR Trials, and for emergency EVRAR a summary of published contemporary practise. Abbreviations: common iliac artery (CIA); external iliac artery (EIA).

In the haemodynamically unstable patient with RAAA, rapid control of progressive haemorrhage at the aortic rupture site during open surgery is often considered paramount. In the setting of endovascular repair, this control may be achieved by swift endograft placement and deployment, or by placement and inflation of a balloon occlusion device in the aorta proximal to the rupture site.^{35,44,72} The reported studies describe the use of both, aorto-uni-iliac (AUI) or aorto-bi-iliac (ABI) devices for EVRAR. Aorto-uni-iliac devices may have the advantage of ease in deployment, and therefore rapid control of haemorrhage, but this approach must be accompanied by a surgical fem-fem crossover bypass graft to provide blood flow to the contralateral lower limb. Aorto-bi-iliac devices, or modular bifurcated grafts, give a better anatomical result but often require more deployment time for cannulation of the contralateral graft limb. However, these are the device of choice for elective EVAR, such that many endovascular specialists are familiar with their capabilities can obtain temporary control by initial proximal deployment of the graft body, delaying release of the contralateral limb until cannulation is imminent or the patient has been stabilized.⁷² Exposure of the femoral arteries for endograft placement under local anaesthetic is advocated by many and can reduce the haemodynamic instability.⁷³ Furthermore, a policy of permissive hypotension to reduce bleeding and prevent re-bleeding from the contained aortic rupture site,^{13,16} is applied in all studies but with wide variation in the lowest tolerated systolic blood pressures (SBP), with most authors considering an un-assisted SBP greater than 70 mm Hg as indicative of stability).^{13,16,40} However, it is notable that even when SBP as low as 50 mm Hg were permitted by

some, without increase in end-organ injury, such as visceral ischaemia.⁴⁰

Patient outcomes after EVRAR

For elective aneurysm repair, EVAR is associated with significant reductions in early complications and mortality.^{11,50} Paradoxically, this early mortality benefit is not sustained at one-year, were the mortality from EVAR is comparable to open repair but at an increased cost.¹¹ Furthermore, high-risk patients deemed unfit for elective open surgical repair, fared no better with EVAR than with best medical management in terms of aneurysm-related or all-cause mortality, suggesting that unfit for open surgery as judged in these studies may be an indicator for reduced life-expectancy.¹⁰ Notable also from the EVAR 2 Trial is that the peri-procedural mortality rate of 9% in these high-risk patients was significantly greater than their lower risk counterparts in EVAR 1 Trial.^{10,11} However, the risk/benefit analysis is different in patients with RAAA as failure to treat means inevitable death and conventional open surgery continues to carry a very significant risk of mortality, 35% to 70%. Endovascular ruptured aneurysm repair (EVRAR) is less invasive, reduces surgical stress, reduces haemodynamic instability, and can be achieved with a local or loco-regional anaesthesia. The only published RCT showed no benefit for EVRAR in terms of mortality or complications.¹⁶ Indeed mortality in both EVRAR and conventional open groups, in a heavily selected population where unstable patients were excluded, was high at 53% compared to many contemporary reports.¹⁶ In the non-randomised studies EVRAR was associated with a low mortality rate, average 17% (range 0–45), compared in some studies to contemporary or historical control groups undergoing open repair, average 34% (range 0–70), for patients with ruptured abdominal aortic aneurysm. Furthermore, this mortality benefit mirrors a procedural-related reduction in blood loss, transfusion requirements, and length of ICU stay (Tables 2–4). These perceived benefits are generally attributed to a reduction in the physiological insult to the patients, as EVRAR obviates the need for laparotomy, exposure and handling of abdominal contents, and aorto-iliac dissection and clamping. In the majority of the studies EVRAR was conducted under general anaesthetic, although it is clear that even AUI device placement with surgical femoro-femoral bypass graft can be achieved under local anaesthetic regimes.⁷³ However, results from selected populations in these non-randomised studies must be interpreted with caution.

Implications for vascular services

A distinct learning curve effect is noted in the studies involving EVRAR, which especially is important as all of these arise from centres with already considerable experience with elective endovascular aneurysm procedures.^{14,25,35,44} The more recent studies show a greater reduction in procedure times, mortality and complication rates.^{18,22,45} As has been seen with elective EVAR practice, advancements in stent-graft design and endovascular techniques have lead to improved outcomes.⁵¹ Re-intervention rates also appear comparable with those seen with elective endovascular repair.^{51,74} However, long-term data are needed in order to truly assess if EVRAR is a durable treatment in relation to endoleak, stent-graft integrity, and late rupture risk. It is clear that the introduction of an EVRAR service has substantial cost implications, in terms of staff, fixed resources and procedure-associated equipment. In order to provide a comprehensive 24-hour service the necessary team of surgeons, radiologists, anaesthetists, radiographers, nurses and technicians need to be available at all times.¹⁵ This may impact on the transferability of this technique beyond specialist centres. These logistical deficiencies were highlighted by the low recruitment rate to the RCT even from a very experienced endovascular centre.¹⁶ Whilst, most units had a wide range of stent-graft stock available to cope with the variable anatomy encountered, the majority of patient can be treated with a small range of devices made available by satisfactory arrangement with a commercial partner. Surveillance protocols in respect to imaging after EVRAR may be similar to elective practice, although the frequency of medium and late complications is as yet unknown and may require a high rate of re-intervention by endovascular or open means, in relation to endoleak, device migration, strut fracture, limb occlusion or late rupture.^{15,70,75}

Study limitations

In the absence of high-quality randomised controlled trials, we are unable to fully evaluate the role of emergency endovascular ruptured abdominal aortic aneurysm repair (EVRAR). Early results from specialist centres show this technique is feasible. Data suggesting reduced morbidity and mortality in selected patients must be interpreted with caution due to the lack of randomisation in these studies. Further trials to evaluate the role of EVRAR in the treatment of RAAA are required. These trials should be methodologically adequate in terms of sample sizes, treatment

standardization and duration of follow up. Clinically-relevant outcomes such as rate of major complications, open-conversion, aneurysm exclusion, endoleak, rupture, and mortality should be assessed. However, accumulating evidence from non-randomised studies, which show significant reductions in mortality in selected patients deemed suitable for endovascular repair, may raise ethical concerns in relation to informed consent and randomisation of these patients to open repair. Large prospective studies are required to validate the acceptable anatomical criteria for EVRAR in RAAA. Furthermore, longitudinal studies are required to assess the long-term durability of this form of treatment in terms of re-intervention rate, open-conversion rate, and rupture-free survival.

Conclusions

Accumulating published data suggests that endovascular ruptured abdominal aortic aneurysm repair (EVRAR) is feasible in selected patients in institutions with experience of endovascular techniques for the treatment of ruptured abdominal aortic aneurysm. Furthermore, in those selected patients EVRAR may also be accompanied by reductions in blood loss, Intensive Care Unit (ICU) stay, and mortality. However, a significant proportion of patients with ruptured abdominal aortic aneurysm remain anatomically unsuitable for contemporary endovascular repair, and whilst relaxation of exclusion criteria may make EVRAR feasible, this is likely to increase device and aneurysm related complications.

Acknowledgements

We the authors would like to thank for their assistance in the preparation of this review the Peripheral Arterial Disease Group, The Cochrane Collaboration. Internal sources of support: Cochrane Fellowship, Research & Development Office, Northern Ireland. Potential conflict of interest: none known.

References

- 1 SCOTT RA, WILSON NM, ASHTON HA, KAY DN. Influence of screening on the incidence of ruptured abdominal aortic aneurysm: 5-year results of a randomized controlled study. *Br J Surg* 1995; **82**:1066–1070.
- 2 SCOTT RA, VARDOLAKI KA, WALKER NM, DAY NE, DUFFY SW, ASHTON HA *et al*. The long-term benefits of a single scan for abdominal aortic aneurysm (AAA) at age 65. *Eur J Vasc Endovasc Surg* 2001;**21**:535–540.
- 3 ASHTON HA, BUXTON MJ, DAY NE, KIM LG, MARTEAU TM, SCOTT RA *et al*. The Multicentre Aneurysm Screening Study (MASS) into the effect of abdominal aortic aneurysm screening on mortality in men: a randomised controlled trial. *Lancet* 2002;**360**:1531–1539.

- 4 FILIPOVIC M, GOLDACRE MJ, ROBERTS SE, YEATES D, DUNCAN ME, COOK-MOZAFFARI P *et al.* Trends in mortality and hospital admission rates for abdominal aortic aneurysm in England and Wales, 1979–1999. *Br J Surg* 2005;**92**:968–975.
- 5 GREENHALGH RM, FORBES JF, FOWKES FG, POWEL JT, RUCKLEY CV, BRADY AR *et al.* Early elective open surgical repair of small abdominal aortic aneurysms is not recommended: results of the UK Small Aneurysm Trial. Steering Committee. *Eur J Vasc Endovasc Surg* 1998;**16**:462–464.
- 6 HUBER TS, HARWARD TR, FLYNN TC, ALBRIGHT JL, SEEGER JM. Operative mortality rates after elective infrarenal aortic reconstructions. *J Vasc Surg* 1995;**22**:287–293.
- 7 KAZMERS A, JACOBS L, PERKINS A, LINDENAUER SM, BATES E. Abdominal aortic aneurysm repair in Veterans Affairs medical centers. *J Vasc Surg* 1996;**23**:191–200.
- 8 BAYLY PJ, MATTHEWS JN, DOBSON PM, PRICE ML, THOMAS DG. In-hospital mortality from abdominal aortic surgery in Great Britain and Ireland: Vascular Anaesthesia Society audit. *Br J Surg* 2001;**88**:687–692.
- 9 NEARY WD, CROW P, FOY C, PRYTHERCH D, HEATHER BP, EARNSHAW JJ *et al.* Comparison of POSSUM scoring and the Hardman Index in selection of patients for repair of ruptured abdominal aortic aneurysm. *Br J Surg* 2003;**90**:421–425.
- 10 Endovascular aneurysm repair and outcome in patients unfit for open repair of abdominal aortic aneurysm (EVAR trial 2): randomised controlled trial. *Lancet* 2005;**365**:2187–2192.
- 11 Endovascular aneurysm repair versus open repair in patients with abdominal aortic aneurysm (EVAR trial 1): randomised controlled trial. *Lancet* 2005;**365**:2179–2186.
- 12 CUYPERS PW, GARDIEN M, BUTH J, PEELS CH, CHARBON JA, HOP WC *et al.* Randomized study comparing cardiac response in endovascular and open abdominal aortic aneurysm repair. *Br J Surg* 2001;**88**:1059–1065.
- 13 PEPPELENBOSCH N, YILMAZ N, VAN MARREWIJK C, BUTH J, CUYPERS P, DUIJM L *et al.* Emergency treatment of acute symptomatic or ruptured abdominal aortic aneurysm. Outcome of a prospective intent-to-treat by EVAR protocol. *Eur J Vasc Endovasc Surg* 2003;**26**:303–310.
- 14 HINCHLIFFE RJ, YUSUF SW, MACIEREWICZ JA, MACSWEENEY ST, WENHAM PW, HOPKINSON BR *et al.* Endovascular repair of ruptured abdominal aortic aneurysm—a challenge to open repair? Results of a single centre experience in 20 patients. *Eur J Vasc Endovasc Surg* 2001;**22**:528–534.
- 15 RESCH T, MALINA M, LINDBLAD B, DIAS NV, SONESSON B, IVANCEV K *et al.* Endovascular repair of ruptured abdominal aortic aneurysms: logistics and short-term results. *J Endovasc Ther* 2003;**10**:440–446.
- 16 HINCHLIFFE RJ, BRUIJSTENS L, MACSWEENEY ST, BRAITHWAITE BD. A randomised trial of endovascular and open surgery for ruptured abdominal aortic aneurysm — results of a pilot study and lessons learned for future studies. *Eur J Vasc Endovasc Surg* 2006;**32**:506–513.
- 17 ALSAC JM, DESGRANGES P, KOBEITER H, BECQUEMIN JP. Emergency endovascular repair for ruptured abdominal aortic aneurysms: feasibility and comparison of early results with conventional open repair. *Eur J Vasc Endovasc Surg* 2005;**30**:632–639.
- 18 ARYA N, MAKAR RR, LAU LL, LOAN W, LEE B, HANNON RJ *et al.* An intention-to-treat by endovascular repair policy may reduce overall mortality in ruptured abdominal aortic aneurysm. *J Vasc Surg* 2006;**44**:467–471.
- 19 BRANDT M, WALLUSCHECK KP, JAHNKE T, GRAW K, CREMER J, MULLER-HULSBECK S *et al.* Endovascular repair of ruptured abdominal aortic aneurysm: feasibility and impact on early outcome. *J Vasc Interv Radiol* 2005;**16**:1309–1312.
- 20 CASTELLI P, CARONNO R, PIFFARETTI G, TOZZI M, LAGANA D, CARRAFIELLO G *et al.* Ruptured abdominal aortic aneurysm: endovascular treatment. *Abdom Imaging* 2005;**30**:263–269.
- 21 DOSS M, MARTENS S, HEMMER W. Emergency endovascular interventions for ruptured thoracic and abdominal aortic aneurysms. *Am Heart J* 2002;**144**:544–548.
- 22 FRANKS S, LLOYD G, FISHWICK G, BOWN M, SAYERS R. Endovascular treatment of ruptured and symptomatic abdominal aortic aneurysms. *Eur J Vasc Endovasc Surg* 2006;**31**:345–350.
- 23 GERASSIMIDIS TS, PAPAZOGLU KO, KAMPARODIS AG, KONSTANTINIDIS K, KARKOS CD, KARAMANOS D *et al.* Endovascular management of ruptured abdominal aortic aneurysms: 6-year experience from a Greek center. *J Vasc Surg* 2005;**42**:615–623.
- 24 GRECO G, EGOROVA N, ANDERSON PL, GELIJNS A, MOSKOWITZ A, NOWYGRAD R *et al.* Outcomes of endovascular treatment of ruptured abdominal aortic aneurysms. *J Vasc Surg* 2006;**43**:453–459.
- 25 GREENBERG RK, SRIVASTAVA SD, OURIEL K, WALDMAN D, IVANCEV K, ILLIG KA *et al.* An endoluminal method of hemorrhage control and repair of ruptured abdominal aortic aneurysms. *J Endovasc Ther* 2000;**7**:1–7.
- 26 HECHELHAMMER L, LACHAT ML, WILDERMUTH S, BETTEX D, MAYER D, PFAMMATTER T *et al.* Midterm outcome of endovascular repair of ruptured abdominal aortic aneurysms. *J Vasc Surg* 2005;**41**:752–757.
- 27 KAPMA MR, VERHOEVEN EL, TIELLIU IF, ZEEBREGTS CJ, PRINS TR, VAN DER HB *et al.* Endovascular treatment of acute abdominal aortic aneurysm with a bifurcated stentgraft. *Eur J Vasc Endovasc Surg* 2005;**29**:510–515.
- 28 LACHAT ML, PFAMMATTER T, WITZKE HJ, BETTEX D, KUNZLI A, WOLFENBERGER U *et al.* Endovascular repair with bifurcated stent-grafts under local anaesthesia to improve outcome of ruptured aortoiliac aneurysms. *Eur J Vasc Endovasc Surg* 2002;**23**:528–536.
- 29 LAGANA D, CARRAFIELLO G, MANGINI M, GIORGIANNI A, CARONNO R, CASTELLI P *et al.* Management and endovascular treatment of symptomatic abdominal aortic aneurysms. *Radiol Med (Torino)* 2006;**111**:959–970.
- 30 LARZON T, LINDGREN R, NORGREN L. Endovascular treatment of ruptured abdominal aortic aneurysms: a shift of the paradigm? *J Endovasc Ther* 2005;**12**:548–555.
- 31 LEE WA, HIRNEISE CM, TAYYARAH M, HUBER TS, SEEGER JM. Impact of endovascular repair on early outcomes of ruptured abdominal aortic aneurysms. *J Vasc Surg* 2004;**40**:211–215.
- 32 LOMBARDI JV, FAIRMAN RM, GOLDEN MA, CARPENTER JP, MITCHELL M, BARKER C *et al.* The utility of commercially available endografts in the treatment of contained ruptured abdominal aortic aneurysm with hemodynamic stability. *J Vasc Surg* 2004;**40**:154–160.
- 33 MEHTA M, DARLING III RC, RODDY SP, FECTEAU S, OZSVATH KJ, KREIENBERG PB *et al.* Factors associated with abdominal compartment syndrome complicating endovascular repair of ruptured abdominal aortic aneurysms. *J Vasc Surg* 2005;**42**:1047–1051.
- 34 OHKI T, VEITH FJ, SANCHEZ LA, CYNAMON J, LIPSITZ EC, WAIN RA *et al.* Endovascular graft repair of ruptured aortoiliac aneurysms. *J Am Coll Surg* 1999;**189**:102–112.
- 35 OHKI T, VEITH FJ. Endovascular grafts and other image-guided catheter-based adjuncts to improve the treatment of ruptured aortoiliac aneurysms. *Ann Surg* 2000;**232**:466–479.
- 36 OREND KH, KOTSIS T, SCHARRE-PAMLER R, KAPFER X, LIEWALD F, GORICH J *et al.* Endovascular repair of aortic rupture due to trauma and aneurysm. *Eur J Vasc Endovasc Surg* 2002;**23**:61–67.
- 37 PEPPELENBOSCH N, CUYPERS PW, VAHL AC, VERMASSSEN F, BUTH J. Emergency endovascular treatment for ruptured abdominal aortic aneurysm and the risk of spinal cord ischemia. *J Vasc Surg* 2005;**42**:608–614.
- 38 REICHAERT M, GEELKERKEN RH, HUISMAN AB, VAN DET RJ, DE SMIT P, VOLKER EP *et al.* Ruptured abdominal aortic aneurysm: endovascular repair is feasible in 40% of patients. *Eur J Vasc Endovasc Surg* 2003;**26**:479–486.
- 39 RUBIN BG, SANCHEZ LA, CHOI ET, SICARD GA. Endoluminal repair of ruptured abdominal aortic aneurysms under local anesthesia: initial experience. *Vasc Endovascular Surg* 2004;**38**:203–207.
- 40 SCHARRE-PAMLER R, KOTSIS T, KAPFER X, GORICH J, SUNDERPLASSMANN L. Endovascular stent-graft repair of ruptured aortic aneurysms. *J Endovasc Ther* 2003;**10**:447–452.
- 41 VADDINENI SK, RUSSO GC, PATTERSON MA, TAYLOR SM, JORDAN Jr WD. Ruptured abdominal aortic aneurysm:

- a retrospective assessment of open versus endovascular repair. *Ann Vasc Surg* 2005;19:782–786.
- 42 VAN SAMBEEK MR, VAN DIJK LC, HENDRIKS JM, VAN GROTEL M, KUIPER JW, PATTYNAMA PM *et al.* Endovascular versus conventional open repair of acute abdominal aortic aneurysm: feasibility and preliminary results. *J Endovasc Ther* 2002;9:443–448.
 - 43 VAN HI, VERMASSEN F, DURIEUX C, RANDON C, DE ROOSE J. Endovascular repair of aortic rupture. *Eur J Vasc Endovasc Surg* 2003;26:311–316.
 - 44 VERHOEVEN EL, PRINS TR, VAN DEN DUNGEN JJ, TIELLIU IF, HULSEBOS RG, VAN SCHILFGAARDE R *et al.* Endovascular repair of acute AAAs under local anesthesia with bifurcated endografts: a feasibility study. *J Endovasc Ther* 2002;9:729–735.
 - 45 VISSER JJ, BOSCH JL, HUNINK MG, VAN DIJK LC, HENDRIKS JM, POLDERMANS D *et al.* Endovascular repair versus open surgery in patients with ruptured abdominal aortic aneurysms: clinical outcomes with 1-year follow-up. *J Vasc Surg* 2006;44:1148–1155.
 - 46 YILMAZ N, PEPELENBOSCH N, CUYPERS PW, TIELBEEK AV, DUIJM LE, BUTH J *et al.* Emergency treatment of symptomatic or ruptured abdominal aortic aneurysms: the role of endovascular repair. *J Endovasc Ther* 2002;9:449–457.
 - 47 GLIMAKER H, HOLMBERG L, ELVIN A, NYBACKA O, ALMGREN B, BJORCK CG *et al.* Natural history of patients with abdominal aortic aneurysm. *Eur J Vasc Surg* 1991;5:125–130.
 - 48 DIMICK JB, STANLEY JC, AXELROD DA, KAZMERS A, HENKE PK, JACOBS LA *et al.* Variation in death rate after abdominal aortic aneurysmectomy in the United States: impact of hospital volume, gender, and age. *Ann Surg* 2002;235:579–585.
 - 49 PARODI JC, PALMAZ JC, BARONE HD. Transfemoral intraluminal graft implantation for abdominal aortic aneurysms. *Ann Vasc Surg* 1991;5:491–499.
 - 50 PRINSEN M, VERHOEVEN EL, BUTH J, CUYPERS PW, VAN SAMBEEK MR, BALM R *et al.* A randomized trial comparing conventional and endovascular repair of abdominal aortic aneurysms. *N Engl J Med* 2004;351:1607–1618.
 - 51 HARRIS PL, BUTH J. An update on the important findings from the EUROSTAR EVAR registry. *Vascular* 2004;12:33–38.
 - 52 FRANSEN GA, VALLABHANENI SR, VAN MARREWIK CJ, LAHEIJ RJ, HARRIS PL, BUTH J *et al.* Rupture of infra-renal aortic aneurysm after endovascular repair: a series from EUROSTAR registry. *Eur J Vasc Endovasc Surg* 2003;26:487–493.
 - 53 BOWN MJ, NICHOLSON ML, BELL PR, SAYERS RD. Cytokines and inflammatory pathways in the pathogenesis of multiple organ failure following abdominal aortic aneurysm repair. *Eur J Vasc Endovasc Surg* 2001;22:485–495.
 - 54 BOWN MJ, NICHOLSON ML, BELL PR, SAYERS RD. The systemic inflammatory response syndrome, organ failure, and mortality after abdominal aortic aneurysm repair. *J Vasc Surg* 2003;37:600–606.
 - 55 HADJIANASTASSIOU VG, TEKKIS PP, GOLDHILL DR, HANDS LJ. Quantification of mortality risk after abdominal aortic aneurysm repair. *Br J Surg* 2005;92:1092–1098.
 - 56 PRYTHERCH DR, SUTTON GL, BOYLE JR. Portsmouth POSSUM models for abdominal aortic aneurysm surgery. *Br J Surg* 2001;88:958–963.
 - 57 HINTERSEHER I, SAEGER HD, KOCH R, BLOOMENTHAL A, OCKERT D, BERGERT H *et al.* Quality of life and long-term results after ruptured abdominal aortic aneurysm. *Eur J Vasc Endovasc Surg* 2004;28:262–269.
 - 58 PEARSON S, HASSEN T, SPARK JL, CABOT J, COWLED P, FITZGIBB R *et al.* Endovascular repair of abdominal aortic aneurysm reduces intraoperative cortisol and perioperative morbidity. *J Vasc Surg* 2005;41:919–925.
 - 59 HINCHLIFFE RJ, HOPKINSON BR. Ruptured abdominal aortic aneurysm. Time for a new approach. *J Cardiovasc Surg (Torino)* 2002;43:345–347.
 - 60 YUSUF SW, WHITAKER SC, CHUTER TA, WENHAM PW, HOPKINSON BR. Emergency endovascular repair of leaking aortic aneurysm. *Lancet* 1994;344:1645.
 - 61 VEITH FJ, BAUM RA, OHKI T, AMOR M, ADISESHIAH M, BLANKENSTEIN JD *et al.* Nature and significance of endoleaks and endotension: summary of opinions expressed at an international conference. *J Vasc Surg* 2002;35:1029–1035.
 - 62 ALSAC JM, KOBEITER H, BECQUEMIN JP, DESGRANGES P. Endovascular repair for ruptured AAA: a literature review. *Acta Chir Belg* 2005;105:134–139.
 - 63 HINCHLIFFE RJ, ALRIC P, ROSE D, OWEN V, DAVIDSON IR, ARMON MP *et al.* Comparison of morphologic features of intact and ruptured aneurysms of infrarenal abdominal aorta. *J Vasc Surg* 2003;38:88–92.
 - 64 ROSE DF, DAVIDSON IR, HINCHLIFFE RJ, WHITAKER SC, GREGSON RH, MACSWEENEY ST *et al.* Anatomical suitability of ruptured abdominal aortic aneurysms for endovascular repair. *J Endovasc Ther* 2003;10:453–457.
 - 65 STERNBERGH III WC, MONEY SR, GREENBERG RK, CHUTER TA. Influence of endograft oversizing on device migration, endoleak, aneurysm shrinkage, and aortic neck dilation: results from the Zenith Multicenter Trial. *J Vasc Surg* 2004;39:20–26.
 - 66 LLOYD GM, BOWN MJ, NORWOOD MG, DEB R, FISHWICK G, BELL PR *et al.* Feasibility of preoperative computer tomography in patients with ruptured abdominal aortic aneurysm: a time-to-death study in patients without operation. *J Vasc Surg* 2004;39:788–791.
 - 67 PEPELENBOSCH N, BUTH J, HARRIS PL, VAN MARREWIK C, FRANSEN G. Diameter of abdominal aortic aneurysm and outcome of endovascular aneurysm repair: does size matter? A report from EUROSTAR. *J Vasc Surg* 2004;39:288–297.
 - 68 WILSON WR, FISHWICK G, SIR PETER RFB, THOMPSON MM. Suitability of ruptured AAA for endovascular repair. *J Endovasc Ther* 2004;11:635–640.
 - 69 VEITH FJ, OHKI T, LIPSITZ EC, SUGGS WD, CYNAMON J. Treatment of ruptured abdominal aneurysms with stent grafts: a new gold standard? *Semin Vasc Surg* 2003;16:171–175.
 - 70 BADGER SA, O'DONNELL ME, MAKAR RR, LOAN W, LEE B, SOONG CV *et al.* Aortic necks of ruptured abdominal aneurysms dilate more than asymptomatic aneurysms after endovascular repair. *J Vasc Surg* 2006;44:244–249.
 - 71 BOYLE JR, GIBBS PJ, KRUGER A, SHEARMAN CP, RAPTSIS S, PHILLIPS MJ *et al.* Existing delays following the presentation of ruptured abdominal aortic aneurysm allow sufficient time to assess patients for endovascular repair. *Eur J Vasc Endovasc Surg* 2005;29:505–509.
 - 72 HINCHLIFFE RJ, BRAITHWAITE BD, HOPKINSON BR. The endovascular management of ruptured abdominal aortic aneurysms. *Eur J Vasc Endovasc Surg* 2003;25:191–201.
 - 73 MORALES JP, IRANI FG, JONES KG, TAYLOR PR, DOURADO R, SABHARWAL T *et al.* Case report: endovascular repair of a ruptured abdominal aortic aneurysm under local anaesthesia. *Br J Radiol* 2005;78:62–64.
 - 74 THOMAS SM, BEARD JD, IRELAND M, AYERS S. Results from the prospective registry of endovascular treatment of abdominal aortic aneurysms (RETA): mid term results to five years. *Eur J Vasc Endovasc Surg* 2005;29:563–570.
 - 75 PRINSEN M, WIXON CL, BUSKENS E, BLANKENSTEIN JD. Surveillance after endovascular aneurysm repair: diagnostics, complications, and associated costs. *Ann Vasc Surg* 2004.

Accepted 3 June 2007

Available online 27 August 2007